

Hyaline Fibromatosis Syndrome

Subjects: Genetics & Heredity

Contributor: Camila Xu

Hyaline fibromatosis syndrome is a disorder in which a clear (hyaline) substance abnormally accumulates in body tissues. This disorder affects many areas of the body, including the skin, joints, bones, and internal organs.

Keywords: genetic conditions

1. Introduction

The severity of the signs and symptoms of hyaline fibromatosis syndrome fall along a spectrum. In more severe cases (previously diagnosed as infantile systemic hyalinosis), signs and symptoms are present at birth or begin within the first few months of life and can be life-threatening. In milder cases (previously diagnosed as juvenile hyaline fibromatosis), signs and symptoms begin in childhood and affect fewer body systems.

One of the main features of hyaline fibromatosis syndrome is the growth of noncancerous masses of tissue (nodules) under the skin, very commonly on the scalp. In more severely affected individuals, nodules also grow in the muscles and internal organs, causing pain and complications. Some severely affected individuals develop a condition called protein-losing enteropathy due to the formation of nodules in their intestines. This condition results in severe diarrhea, failure to gain weight and grow at the expected rate, and general wasting and weight loss (cachexia).

Another common feature of hyaline fibromatosis syndrome is painful skin bumps that frequently appear on the hands, neck, scalp, ears, and nose. They can also develop in joint creases and the genital region. These skin bumps are described as white or pink and pearly. They may be large or small and often increase in number over time.

In some affected individuals, especially those with more severe signs and symptoms, the skin covering joints, such as the ankles, wrists, elbows, and finger joints, is unusually dark (hyperpigmented). Hyaline fibromatosis syndrome is also characterized by overgrowth of the gums (gingival hypertrophy), and some affected individuals have thickened skin.

Joint stiffness and pain are common in hyaline fibromatosis syndrome, and many affected individuals develop joint deformities called contractures that limit movement. By adulthood, some people with the condition require a wheelchair for mobility. Bone abnormalities can also occur in hyaline fibromatosis syndrome.

Although individuals with hyaline fibromatosis syndrome have severe physical limitations, mental development is typically normal. People with milder signs and symptoms live into adulthood, while the most severely affected individuals often do not survive beyond early childhood due to chronic diarrhea and recurrent infections.

2. Frequency

Hyaline fibromatosis syndrome is a rare condition. Its prevalence is unknown.

3. Causes

Hyaline fibromatosis syndrome is caused by mutations in a gene called *ANTXR2*. This gene provides instructions for making a protein that is found at the surface of many types of cells. The *ANTXR2* protein is believed to interact with components of the extracellular matrix, which is the lattice of proteins and other molecules outside the cell. This matrix strengthens and supports connective tissues, such as skin, bone, cartilage, tendons, and ligaments. The *ANTXR2* protein may play a role in the structure of the extracellular matrix. The nature of the hyaline substance that builds up in hyaline fibromatosis syndrome is unknown, but it likely contains extracellular matrix proteins, among other materials.

Mutations in the *ANTXR2* gene are thought to result in production of an ANTXR2 protein that is unable to get to the surface of cells or that has impaired ability to interact with extracellular matrix components. It is unclear what effect these mutations have in cells and tissues. Researchers suspect that gene mutations disrupt the formation of the extracellular matrix, allowing a hyaline substance to leak through and build up in various body tissues. Alternatively, the mutations could impair the breakdown of excess extracellular matrix proteins, which then accumulate in tissues and lead to the signs and symptoms of hyaline fibromatosis syndrome.

Researchers are unsure why the severity of hyaline fibromatosis syndrome varies among affected individuals. Some studies have indicated that the severity of the condition may be linked to where in the gene the mutation occurs.

3.1. The gene associated with Hyaline fibromatosis syndrome

- ANTXR2

4. Inheritance

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

5. Other Names for This Condition

- inherited systemic hyalinosis
- mollusum fibrosum
- Murray syndrome
- Puretic syndrome

References

1. Bürgi J, Kunz B, Abrami L, Deuquet J, Piersigilli A, Scholl-Bürgi S, Lausch E, Unger S, Superti-Furga A, Bonaldo P, van der Goot FG. CMG2/ANTXR2 regulates extracellular collagen VI which accumulates in hyaline fibromatosis syndrome. *Nat Commun*. 2017 Jun 12;8:15861. doi: 10.1038/ncomms15861.
2. Denadai R, Raposo-Amaral CE, Bertola D, Kim C, Alonso N, Hart T, Han S, Stelini RF, Buzzo CL, Raposo-Amaral CA, Hart PS. Identification of 2 novel ANTXR2 mutations in patients with hyaline fibromatosis syndrome and proposal of a modified grading system. *Am J Med Genet A*. 2012 Apr;158A(4):732-42. doi:10.1002/ajmg.a.35228.
3. Deuquet J, Abrami L, Difeo A, Ramirez MC, Martignetti JA, van der Goot FG. Systemic hyalinosis mutations in the CMG2 ectodomain leading to loss of function through retention in the endoplasmic reticulum. *Hum Mutat*. 2009 Apr;30(4):583-9. doi: 10.1002/humu.20872.
4. Deuquet J, Lausch E, Superti-Furga A, van der Goot FG. The dark sides of capillary morphogenesis gene 2. *EMBO J*. 2012 Jan 4;31(1):3-13. doi:10.1038/emboj.2011.442.
5. El-Kamah GY, Fong K, El-Ruby M, Afifi HH, Clements SE, Lai-Cheong JE, Amr K, El-Darouti M, McGrath JA. Spectrum of mutations in the ANTXR2 (CMG2) gene in infantile systemic hyalinosis and juvenile hyaline fibromatosis. *Br J Dermatol*. 2010 Jul;163(1):213-5. doi: 10.1111/j.1365-2133.2010.09769.x.
6. Tanaka K, Ebihara T, Kusubata M, Adachi E, Arai M, Kawaguchi N, Utsunomiya J, Miki Y, Hiramoto M, Hattori S, Irie S. Abnormal collagen deposition in fibromas from patient with juvenile hyaline fibromatosis. *J Dermatol Sci*. 2009 Sep;55(3):197-200. doi: 10.1016/j.jdermsci.2009.06.005.
7. Yan SE, Lemmin T, Salvi S, Lausch E, Superti-Furga A, Rokicki D, Dal Peraro M, van der Goot FG. In-depth analysis of hyaline fibromatosis syndrome frameshift mutations at the same site reveal the necessity of personalized therapy. *Hum Mutat*. 2013 Jul;34(7):1005-17. doi: 10.1002/humu.22324.

