# **Neurofibromatosis Type 2**

Subjects: Genetics & Heredity Contributor: Rita Xu

Neurofibromatosis type 2 is a disorder characterized by the growth of noncancerous tumors in the nervous system.

Keywords: genetic conditions

## 1. Introduction

The most common tumors associated with neurofibromatosis type 2 are called vestibular schwannomas or acoustic neuromas. These growths develop along the nerve that carries information from the inner ear to the brain (the auditory nerve). Tumors that occur on other nerves are also commonly found with this condition.

The signs and symptoms of neurofibromatosis type 2 usually appear during adolescence or in a person's early twenties, although they can begin at any age. The most frequent early symptoms of vestibular schwannomas are hearing loss, ringing in the ears (tinnitus), and problems with balance. In most cases, these tumors occur in both ears by age 30. If tumors develop elsewhere in the nervous system, signs and symptoms vary according to their location. Complications of tumor growth can include changes in vision, numbness or weakness in the arms or legs, and fluid buildup in the brain. Some people with neurofibromatosis type 2 also develop clouding of the lens (cataracts) in one or both eyes, often beginning in childhood.

### 2. Frequency

Neurofibromatosis type 2 has an estimated incidence of 1 in 33,000 people worldwide.

# 3. Causes

Mutations in the *NF2* gene cause neurofibromatosis type 2. The *NF2* gene provides instructions for making a protein called merlin (also known as schwannomin). This protein is produced in the nervous system, particularly in Schwann cells, which surround and insulate nerve cells (neurons) in the brain and spinal cord. Merlin acts as a tumor suppressor, which means that it keeps cells from growing and dividing too rapidly or in an uncontrolled way. Although its exact function is unknown, this protein is likely also involved in controlling cell movement, cell shape, and communication between cells. Mutations in the *NF2* gene lead to the production of a nonfunctional version of the merlin protein that cannot regulate the growth and division of cells. Research suggests that the loss of merlin allows cells, especially Schwann cells, to multiply too frequently and form the tumors characteristic of neurofibromatosis type 2.

#### 3.1. The Gene Associated with Neurofibromatosis Type 2

• NF2

### 4. Inheritance

Neurofibromatosis type 2 is considered to have an autosomal dominant pattern of inheritance. People with this condition are born with one mutated copy of the *NF2* gene in each cell. In about half of cases, the altered gene is inherited from an affected parent. The remaining cases result from new mutations in the *NF2* gene and occur in people with no history of the disorder in their family.

Unlike most other autosomal dominant conditions, in which one altered copy of a gene in each cell is sufficient to cause the disorder, two copies of the *NF2* gene must be altered to trigger tumor formation in neurofibromatosis type 2. A mutation in the second copy of the *NF2* gene occurs in Schwann cells or other cells in the nervous system during a person's lifetime. Almost everyone who is born with one *NF2* mutation acquires a second mutation (known as a somatic mutation) in these cells and develops the tumors characteristic of neurofibromatosis type 2.

# 5. Other Names for This Condition

- BANF
- bilateral acoustic neurofibromatosis
- central neurofibromatosis
- familial acoustic neuromas
- neurofibromatosis 2
- neurofibromatosis type II
- NF2
- schwannoma, acoustic, bilateral

#### References

- 1. Asthagiri AR, Parry DM, Butman JA, Kim HJ, Tsilou ET, Zhuang Z, Lonser RR.Neurofibromatosis type 2. Lancet. 2009 Jun 6;373(9679):1974-86. doi:10.1016/S0140-6736(09)60259-2.
- Evans DG, Howard E, Giblin C, Clancy T, Spencer H, Huson SM, Lalloo F. Birthincidence and prevalence of tumorprone syndromes: estimates from a UK familygenetic register service. Am J Med Genet A. 2010 Feb;152A(2):327-32. doi:10.1002/ajmg.a.33139.
- 3. Evans DG. Neurofibromatosis 2. 1998 Oct 14 [updated 2018 Mar 15]. In: Adam MP,Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors.GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle;1993-2020. Available from http://www.ncbi.nlm.nih.gov/books/NBK1201/
- 4. Evans DG. Neurofibromatosis type 2 (NF2): a clinical and molecular review.Orphanet J Rare Dis. 2009 Jun 19;4:16. doi: 10.1186/1750-1172-4-16. Review.
- Evans GR, Lloyd SKW, Ramsden RT. Neurofibromatosis type 2. AdvOtorhinolaryngol. 2011;70:91-98. doi: 10.1159/000322482.Review.
- 6. Hoa M, Slattery WH 3rd. Neurofibromatosis 2. Otolaryngol Clin North Am. 2012Apr;45(2):315-32, viii. doi: 10.1016/j.otc.2011.12.005. Review.
- Ruggieri M, Iannetti P, Polizzi A, La Mantia I, Spalice A, Giliberto O, Platania N, Gabriele AL, Albanese V, Pavone L. Earliest clinical manifestations and natural history of neurofibromatosis type 2 (NF2) in childhood: a study of 24patients. Neuropediatrics. 2005 Feb;36(1):21-34.

Retrieved from https://encyclopedia.pub/entry/history/show/11802